



NEW STABILITY INDICATING HIGH PERFORMANCE LIQUID CHROMATOGRAPHY METHOD FOR THE DETERMINATION OF NIRMATRELVIR AND RITONAVIR IN BULK AND TABLET DOSAGE FORM

Arthi¹, Dr. S. Sridar², Dr. S Sudhakar³

¹M. Pharmacy, Department of Pharmaceutical Analysis, Malla Reddy College of Pharmacy, Maisammaguda, Secunderabad, Hyderabad, 500100, Medchal District, Secunderabad, Hyderabad, 500100, Medchal District

²Professor, Department of Pharmaceutical Chemistry, Malla Reddy College of Pharmacy, Maisammaguda, Secunderabad, Hyderabad, 500100, Medchal District, Secunderabad, Hyderabad, 500100, Medchal District

³Principal and Professor, Malla Reddy College of Pharmacy, Maisammaguda, Secunderabad, Hyderabad, 500100, Medchal District, Secunderabad, Hyderabad, 500100, Medchal District

Received: 20-12-2024 / Revised Accepted: 24-12-2024 / Published: 02-01-2025

ABSTRACT:

Nirmatrelvir and Ritonavir are prescribed to treat COVID-19. An cheap HPLC approach was developed and is utilized for validation. KH₂PO₄ and MeCN are combined in an Agilent C18 150x4.6mm, 5m column at a flow rate of 1 ml/min at a temperature of 30 oC using a 60:40 v/v ratio. Its length was 215.0 nm. Nirmatrelvir had a holding time of 2.243 hours, whereas Ritonavir took 2.815 hours. It was discovered that the %RSD was 0.5% and 0.5%. The Nirmatrelvir regression equation was: $y = 58409x + 2794$. It was also discovered that ritonavir was $y = 59191x + 774.23$. Additionally, the LOD and LOQ values were discovered, and all tests passed and met ICH criteria

Key Words Nirmatrelvir, Ritonavir, Rp HPLC, Validation, Method Development.

INTRODUCTION

Nirmatrelvir and ritonavir are antiviral agents co-formulated under the brand name Paxlovid, developed by Pfizer for the treatment of COVID-19. Nirmatrelvir is a protease inhibitor targeting the SARS-CoV-2 main protease (Mpro), essential for viral replication. Ritonavir, originally developed as an HIV protease inhibitor, functions here primarily to inhibit cytochrome P450 3A4 (CYP3A4), thereby increasing nirmatrelvir's plasma concentration and prolonging its therapeutic effect.¹

Clinical trials have demonstrated the efficacy of nirmatrelvir-ritonavir in reducing the risk of COVID-19-related hospitalization or death. A pivotal study reported an 89% reduction in such risks among high-risk, non-hospitalized adults treated within five days of symptom onset.²

The U.S. Food and Drug Administration (FDA) granted emergency use authorization for Paxlovid in December 2021, followed by full approval in May 2023 for the treatment of mild-to-moderate COVID-19 in adults at high risk for progression to severe disease.³

Subsequent studies have reinforced the effectiveness of nirmatrelvir-ritonavir in diverse populations, including vaccinated individuals and those with prior COVID-19 infections. Research indicates significant reductions in hospitalization and mortality rates, underscoring its role in the therapeutic landscape of COVID-19.⁴

However, recent investigations into the use of nirmatrelvir-ritonavir for treating post-acute sequelae of SARS-CoV-2 infection (PASC), commonly known as Long COVID, have not demonstrated significant improvements in symptoms compared to placebo.⁵

It is important to note that nirmatrelvir-ritonavir can interact with various medications, potentially leading to serious side effects. Healthcare providers should thoroughly review a patient's medication regimen to manage potential drug interactions effectively.⁶

In summary, the combination of nirmatrelvir and ritonavir represents a significant advancement in the treatment of COVID-19, offering substantial benefits in reducing severe outcomes in high-risk populations. Ongoing research continues to explore its full therapeutic potential and optimal use in various clinical scenarios.⁷

Address for Correspondence: Arthi, M. Pharmacy, Department of Pharmaceutical Analysis, Malla Reddy College of Pharmacy, Maisammaguda, Secunderabad, Hyderabad, 500100, Medchal Dist.; **E-Mail:** arthifnu0@gmail.com

How to Cite this Article: Arthi, New Stability Indicating High Performance Liquid Chromatography Method For The Determination Of Nirmatrelvir And Ritonavir In Bulk And Tablet Dosage Form. World J Pharm Sci 2025; 13(01): 61-68; <https://doi.org/10.54037/WJPS.2022.100905>

Copyright: 2022@ The Author(s). This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License (CC BY-NC-SA), which allows re-users to distribute, remix, adapt, and build upon the material in any medium or format for noncommercial purposes only, and only so long as attribution is given to the creator. If you remix, adapt, or build upon the material, you must license the modified material under identical terms.

Background: Nirmatrelvir is oral protease inhibitor with emergency use authorization for the treatment of mild-to-moderate COVID-19. It is chemically known as (1R,2S,5S)-N-[(1S)-1-cyano-2-[(3S)-2-oxopyrrolidin-3-yl]ethyl]-3-[(2S)-3,3-dimethyl-2-(2,2,2-trifluoroacetamido)butanoyl]-6,6-dimethyl-3-azabicyclo[3.1.0]hexane-2-carboxamide⁸ and Ritonavir is an HIV protease inhibitor chemically known as (1,3-thiazol-5-yl)methyl N-[(2S,3S,5S)-3-hydroxy-5-[(2S)-3-methyl-2-[[methyl({[2-(propan-2-yl)-1,3-thiazol-4-yl]methyl})carbamoyl]amino]butanamido]-1,6-diphenylhexan-2-yl]carbamate⁹




Figure 1: structure of Nirmatrelvir




Figure 2: Structure of Ritonavir

Extensive literature research has unearthed a multitude of recorded analytical procedures, including the discovery of more economically efficient ways. Nevertheless, there is currently no documented approach for calculating stability studies. Hence, a reliable and cost-effective approach is suggested for assessing the stability of Nirmatrelvir, Ritonavir, and their medicinal dose form using RP-HPLC. ¹⁰⁻¹⁸ must be validated and developed as per ICH guidelines.

Materials and Methods: Nirmatrelvir and Ritonavir pure pharmaceuticals (API) are available as gift samples from Spectrum Pharma Research Solution. The chemicals and buffers used in this estimation were supplied by Rankem, an Indian source.

Instrumentation: The development and method validation were conducted using a WATERS HPLC, specifically the model 2695 SYSTEM, equipped with a Photo diode array detector. The system also included an automated sample injector and the Empower 2 software.

Objective: In order to fulfill ICH standards, we need to design and test an HPLC technique that can detect Ritonavir and Nirmatrelvir in pharmaceutical formulations at the same time.

Table 1 : Chromatographic Conditions

Mobile phase	Acetonitrile: OPA (40:60 v/v)
Flow rate	1.0 ml/min
Column	Agilent C18 Column, 5 μ m, 4.6 x 150 mm
Detector wave length	215 nm
Column temperature	26°C
Injection volume	10 μ L
Run time	10.0 min




Figure 3: Optimized Chromatogram

Solution of Standard: 7.5 mg of Nirmatrelvir & 5 mg of Ritonavir of API drug was weighed and added in a separate 50ml vial and it was filled 3/4th volume with the Dil. And then it was under sonication for 10-15 min till

the drug dissolves. (Nirmatrelvir- 150µg/ml, Ritonavir- 100µg/ml) from stock 1ml was taken and added in a 10ml vf and madeup till mark with diluent. (Nirmatrelvir- 15µg/ml, Ritonavir- 10µg/ml)

Sample preparation: Ten tabs takes as mean weight of dose determines to similar to 150 mg and 100 mg. Was acquired Subsequently, 20 ml of acetonitrile was introduced, subjected to sonication for 25 minutes, and adjusted to the mark to achieve concentrations of 1100 and 500 µg/ml. The sample was centrifuges till 20minutes. The sediment subsequently taken and filter using 0.45µm filters. (Nirmatrelvir-300µg/ml, Ritonavir-200µg/ml). from it 1 ml solution was added in a 10ml vf and madeup till mark with diluent. (Nirmatrelvir-15µg/ml, Ritonavir- 10µg/ml)

System suitability parameters: The system suitability parameters were determined by preparing standard solutions of Nirmatrelvir (15ppm) and Ritonavir (10ppm) and the solutions were injected six times and the parameters like peak tailing, resolution and USP plate count were determined.

Specificity: Checking of the interference in the optimized method. We should not find interfering peaks in blank and placebo at retention times of these drugs in this method. So, this method was said to be specific.

Table 2: System suitability results

S.no	Nirmatrelvir			Ritonavir				
	Inj	RT (min)	USP Plate Count	Tailing	RT (min)	USP Plate Count	Tailing	Resolution
1		2.254	4121	1.29	2.805	5264	1.04	3.5
2		2.256	4233	1.29	2.807	5484	1.07	3.6
3		2.257	4246	1.27	2.810	5575	1.06	3.7
4		2.258	4422	1.28	2.813	5241	1.03	3.6
5		2.261	4312	1.27	2.813	5242	1.05	3.7
6		2.261	4295	1.25	2.814	5339	1.03	3.6




Figure 4: system suitability Chromatogram

Table 3: Specificity data

Sample name	Retention time(mins)	Area	Plate count	Resolution
Nirmatrelvir	2.243	249964	1.31	
Ritonavir	2.815	120306	1.05	4.0




Figure 5: Blank

Linearity:

Calibration data and regression data in table 4 and calibration curve in figure 6, 7.

Table 4: Calibration data of Nirmatrelvir and Ritonavir

S. no	Nirmatrelvir		Ritonavir	
	Conc (µg/mL)	Peak area	Conc(µg/mL)	Peak area
1	0	0	0	0
2	3.75	226737	2.5	144621
3	7.5	439995	5	302858
4	11.25	653845	7.5	445501
5	15	881916	10	589965
6	18.75	1105835	12.5	744495
7	22.5	1310944	15	885503
Concentration range	3.75-22.5		2.5-15	
Regression Equation	y = 58409x + 2794		y = 59191x + 774.23	
Co-relation	0.9999		0.9999	
LOD	0.01		0.04	
LOQ	0.04		0.12	




Figure 6: Calibration curve of Nirmatrelvir




Figure 7: Calibration curve of Ritonavir

Accuracy:

Recovery data shown in table 5

Table 5: recovery data of Nirmatrelvir and Ritonavir

% Level	Nirmatrelvir			Ritonavir		
	Amount Spiked (µg/mL)	Amount recovered (µg/mL)	% Recovery	Amount Spiked (µg/mL)	Amount recovered (µg/mL)	% Recovery
50%	7.5	7.42	98.89	5	4.95	99.04
	7.5	7.44	99.26	5	4.96	99.18
	7.5	7.37	98.24	5	4.95	99.09
100%	15	15.09	100.58	10	9.98	99.84
	15	15.08	100.52	10	10.08	100.85
	15	15.05	100.35	10	9.96	99.64
150%	22.5	22.60	100.43	15	14.82	98.78
	22.5	22.31	99.16	15	14.95	99.67
	22.5	22.28	99.01	15	14.80	98.66
% recovery	99.60			99.67		

System precision was performed and the data was shown in table 6

Table 6: System precision of Nirmatrelvir and Ritonavir

	Nirmatrelvir	Ritonavir
1.	883258	596303
2.	885514	594941
3.	879875	596023
4.	878862	596358
5.	880123	589684
6.	873177	590891
Avg	880135	594033
S.D	4214.0	2970.8
%RSD	0.5	0.5

The % RSD for the peak areas of Nirmatrelvir and Ritonavir obtained from six replicate injections of standard solution was within the limit.

Method Precision: The precision of the method was determined by analyzing a sample of Nirmatrelvir and Ritonavir and shown in table 7

Table 7: method Precision

	Nirmatrelvir	Ritonavir
1.	879785	593298
2.	876257	589741
3.	874256	590125
4.	872514	587060
5.	876253	587640
6.	872141	586872
Avg	875201	589123
S.D	2853.5	2463.3
%RSD	0.3	0.4

From the above results, the % RSD of method precision study was within the limit for Nirmatrelvir and Ritonavir.

Robustness: Robustness conditions like Flow minus (0.9ml/min), Flow plus (1.1ml/min), mobile phase minus (55B:45A), mobile phase plus (65B:35A), temperature minus (25°C) and temperature plus(35°C) was maintained and samples were injected in duplicate manner. System suitability parameters were not much affected and all the parameters were passed. %RSD was within the limit.

Table 8: Robustness data for Nirmatrelvir and Ritonavir.

Condition	%R.S. D of Nirmatrelvir	%R.S. D Ritonavir
F - 0.9ml/min	0.2	0.2
F + 1.1ml/min	0.4	0.5
M.P - 55B:45A	0.4	0.5
M.P + 65B:35A	0.2	0.1
T - 25°C	0.1	0.3
T + 35°C	0.3	0.7

Force Degradation Studies: table 9 shows degradation conditions and the obtained degraded data in table 10 and purity plot chromatogram in figure 8,9

Table 9: degradation conditions

Stress condition	Solvent	Temp(0C)	Exposed time
Acid	2N HCL	600c	30 mins
Base	2N NAOH	600c	30 mins
Oxidation	20% H2O2	600c	30 mins
Thermal	Diluent	1050c	6 hours
Photolytic	Diluent	-	-
Hydrolytic	Water	600c	-

Table 10: degradation data

Type of degradation	Nirmatrelvir			Ritonavir		
	area	%recovered	% degraded	area	%recovered	% degraded
Acid	842149	95.49	4.51	559308	93.97	6.03
Base	846710	96.01	3.99	559710	94.03	5.97
Peroxide	849579	96.34	3.66	562908	94.57	5.43
Thermal	859270	97.43	2.57	569041	95.60	4.40
Uv	865864	98.18	1.82	579732	97.40	2.60
Water	875092	98.18	1.82	591049	99.30	0.70




Figure 8: Purity plots for Acid Condition for Nirmatrelvir




Figure 9: Purity plots for Acid Condition for Ritonavir

Assay: Paxlovid Tablet, bearing the label claim Nirmatrelvir 300mg, Ritonavir 300mg. Assay was performed with the above formulation. Average % Assay for Nirmatrelvir and Ritonavir obtained was 99.24% and 98.97% respectively

Table 11: assay data

S.no	Nirmatrelvir			Ritonavir		
	Std Area	Sample area	% Assay	Std Area	Sample area	% Assay
1	883258	879785	99.76	596303	593298	99.68
2	885514	876257	99.36	594941	589741	99.08
3	879875	874256	99.13	596023	590125	99.14
4	878862	872514	98.94	596358	587060	98.63
5	880123	876253	99.36	589684	587640	98.73
6	873177	872141	98.89	590891	586872	98.60
Avg	880135	875201	99.24	594033	589123	98.97
Std ev	4214.0	2853.5	0.32	2970.8	2463.3	0.41
%RSD	0.3	0.3	0.3	0.5	0.3	0.3

Assay was calculated by the formula:

		AT	WS	1	100	10	P	FV	
	% Assay =	X	X	X	X	X	X	X	X 100
		AS	100	10	1	1	100	L.C	
AT		Average Peak area of sample in test solution							
AS		Mean peak area of sample in standard solution							
WS		Weight of drug working standard taken in mg							
P		Assay of drug working standard in % on dried basis							
L.C		Label Claim							

Figure 10 Formula

CONCLUSION:

The findings of the study will be very helpful in evaluating the quality of reasonably priced drugs that contain Ritonavir and Nirmatrelvir. This might be as a result of the study's straightforward sample preparation method, which required little mobile phase and a brief analytical period. The results of evaluating two medications combined in a single dosage demonstrated that the recently created analytical technique was almost entirely successful.

ACKNOWLEDGEMENT:

The authors are thankful to, Department of Pharmaceutical Chemistry, Malla Reddy College of Pharmacy, Affiliated to Osmania University, India and Spectrum Pharma Research Solutions, Hyderabad, Telangana, India.

REFERENCES:

1. Lamb, Y. N. (2022). Nirmatrelvir Plus Ritonavir: First Approval. *Drugs*.
2. Hammond, J., et al. (2022). Oral Nirmatrelvir for High-Risk, Nonhospitalized Adults with Covid-19. *The New England Journal of Medicine*.
3. https://en.wikipedia.org/wiki/Nirmatrelvir/ritonavir?utm_source=chatgpt.com
4. Nirmatrelvir for Vaccinated or Unvaccinated Adult Outpatients with Covid-19. *The New England Journal of Medicine*
5. https://time.com/6986441/paxlovid-does-not-treat-long-covid/?utm_source=chatgpt.com
6. https://medlineplus.gov/druginfo/meds/a622005.html?utm_source=chatgpt.com
7. Heidi Leister-Tebbe et al, Alleviation of Coronavirus Disease 2019 Symptoms and Reduction in Healthcare Utilization Among High-risk Patients Treated With Nirmatrelvir/Ritonavir (NMV/R): A Phase 3 Randomized Trial, *Clinical Infectious Diseases*, ciae551, 2024, <https://doi.org/10.1093/cid/ciae551>
8. <https://go.drugbank.com/drugs/DB16691>
9. <https://go.drugbank.com/drugs/DB00503>
10. Palaniappan Ilayaraja et al., Novel stability indicating HPLC method for the quantification of Nirmatrelvir in bulk drugs, *Elsevier*, Volume 196, January 2024, 109707
11. S. Pallavi et al., Development and validation of a new RP-UPLC method for the simultaneous estimation of nirmatrelvir and ritonavir in bulk and copacked tablet dosage forms, *RJPT*, Vol 16, Issue 9, 2023
12. Mohamed S Imam et al., Adjusted green HPLC determination of nirmatrelvir and ritonavir in the new FDA approved co-packaged pharmaceutical dosage using supported computational calculations, *Sci Rep*, 2023 Jan 4;13(1):137
13. Santosh Kumar Gandhi et al., Development and Validation of RP-HPLC Method for the simultaneous determination of Nirmatrelvir and Ritonavir in bulk and pharmaceutical formulation, *Research Journal of Chemistry and Environment* 27(4):120-127, March 2023
14. Joris Guyon et al., A UPLC-MS/MS Method for Plasma Biological Monitoring of Nirmatrelvir and Ritonavir in the Context of SARS-CoV-2 Infection and Application to a Case, *J Am Soc Mass Spectrom*, 2022 Oct 5;33(10):1975-1981.
15. Srinivas Reddy Jitta et al., Development and Validation of RP-HPLC Method for Quantification of Total, Free and Entrapped Ritonavir in Lipid Nanocarriers and Drug content of Film Coated Fixed Dose Formulation, *Indian Journal of Pharmaceutical Education and Research*, 2022; 56(3s):s547-s558
16. Avinash Birajdar et al., Development and Validation of RP-HPLC method for the Estimation of Ritonavir in API and tablet Formulation, *RJPT*, 14(10): October 2021
17. Andrew J Ocque et al., Development and validation of a UPLC-MS/MS method for the simultaneous determination of Paritaprevir And Ritonavir In Rat Liver, *Bioanalysis* Vol. 8, No. 13 2016.
18. Veera Venkata Satyanarayana Peruri et al., A RP-HPLC Method for the Estimation of Ritonavir in Pharmaceutical dosage forms, *Journal of Pharmacy Research* 2011,4(9),3049-3051.