



Ameliorating Sodium Chloride Stress by Calcium Chloride in *Pennisetum typhoides* Stapf et Hubb

P. Gobinathan*, P. V. Murali

Department of Botany, T.B.M.L. College, Porayar – 609 307, Nagapattinam Dt., Tamil Nadu, India*
Stress Physiology Lab, Department of Botany, Annamalai University, Annamalainagar- 608 002, Tamil Nadu, India

Received: 24-05-2015 / Revised: 29-06-2015 / Accepted: 16-07-2015

ABSTRACT

A pot culture experiment was carried out in *Pennisetum typhoides* Stapf et Hubb. Plants in order to study the effect of calcium chloride (CaCl_2) as an agent ameliorating sodium chloride (NaCl) stress. The plants were raised in pots and salinity stress was imposed by 100 mM of NaCl . The ameliorating effect of CaCl_2 was studied by giving 5mM CaCl_2 to the stressed plants. NaCl stressed plants showed decreased proteins, total sugars, polyphenol oxidase and catalase activity with increased free amino acids, proline and peroxidase activities. When CaCl_2 was combined with NaCl , the CaCl_2 altered overall plant metabolism and paved the way for partial amelioration of the oxidative stress caused by salinity.

Key words: antioxidant enzymes, calcium chloride, *Pennisetum typhoides* Stapf et Hubb, proline, protein, sodium chloride.

Abbreviations: POX – peroxidase activity, PPO – polyphenol oxidase activity, CAT – catalase, DAS – days after sowing.

INTRODUCTION

Salt accumulation in irrigated soils is one of the main factors that diminish crop productivity, since most plants are not halophytic (Hoshida et al. 2000). Salt stress induces various biochemical and physiological responses in plants and affects almost all plant processes (Nemoto and Sasakuma 2002). Plants are subjected to several environmental stresses that adversely affect growth, metabolism and yield (Lawlor 2002). Stresses associated with temperature, salinity and drought singly or in combination are likely to enhance the severity of problems to which plants will be exposed in the coming decades (Duncan 2000). Salinity affects plant growth and germination in a variety of ways i.e., reducing water uptake and nutrient availability, causing toxic accumulation of sodium chloride and other ions. Salinity induces water deficit even in well-watered soils by decreasing the osmotic potential of soil solutes, thus making it difficult for roots to extract water from their surrounding media (Sairam and Srivastava 2002). High concentrations of salt resulting from natural processes or disarrangement in irrigated agriculture result in

inhibition of plant growth and yield (Demiral and Turkan 2006). Salinity is considered a significant factor affecting crop production and agricultural sustainability in arid and semi-arid region of the world, reducing the value and productivity of the affected land (Flowers and Yeo 1992; Munns 1993). The major effect of salinity is the inhibition of crop growth by the reduced hormone delivery from roots to leaves (Azooz et al. 2004). Excessive sodium (Na^+) inhibits the growth of many salt-sensitive plants and glycophytes, which include most crop plants. The typical first response of all plants to salt stress is osmotic adjustment. Compatible solute accumulation in the cytoplasm is considered a mechanism to contribute salt tolerance (Jaleel et al. 2007a). Salinity is considered a major factor in limiting plant growth and crop productivity and salinization of irrigated and surrounding areas in the arid tropics and subtropics has not been diminished, it continues to increase in arid and semi arid region (Teixeira and Pereira 2007).

Adverse effects of salinity on crop growth stem from two characteristics. (1) The increased osmotic potential of the soil solution with salinity makes the

Corresponding Author Address: Dr. P.Gobinathan, Assistant Professor & Head, Department of Botany, Porayar-609 307, Nagapattinam Dt., Tamilnadu, India

water in the soil less available for plants and (2) specific effects of some elements (Na, Cl, B, etc...) present in excess concentration (Yamaguchi and Blumwald 2005; Munus 2005). High salinity is known to cause both hyper ionic and hyper osmotic effects in plants, leading to membrane disorganization and metabolic toxicity (Hasegawa et al. 2000).

An important consequence of salinity stress in plants is the excessive generation of ROS such as the superoxide anion (O_2^-), H_2O_2 and the hydroxyl radicals, particularly in chloroplast and mitochondria (Mittler 2002). In order to survive under stress conditions, plants are equipped with oxygen radical detoxifying enzymes such as superoxide dismutase, ascorbate peroxidase, catalase (and antioxidant molecules like ascorbic acid), α -tocopherol and reduced glutathione (Jaleel et al. 2007b). Antioxidant mechanisms may provide a strategy to enhance salt tolerance in plants.

In plants, calcium ions are the second messenger coupling physiological responses to external and developmental signals (Reddy and Reddy 2004). One possible approach to reducing the effect of salinity on plant productivity is through the addition of calcium supplements to irrigation water (Sohan et al. 1999). Supplementing the medium with Ca^{2+} alleviates growth inhibition by salt in glycophyte plants. Ca^{2+} sustains K^+ transport and K^+/Na^+ selectivity in Na^+ challenged plants. The interaction of Na^+ and Ca^{2+} on plant growth and ion relations is well established (Rengel 1992). The effect of salt stress on nutrition is particularly interesting because Ca^{2+} is one of the important factors involved in the resistance of plants to salt stress. Calcium deficiency symptoms generally arise from differences in its allocation in the growing regions of plants (Greenway and Munns 1980). However effectiveness in alleviating the toxic effect of Na^+ depends on the Ca^{2+} and Na^+ concentration and on the species (Grattan and Grieve 1999). Calcium is known to increase salinity tolerance in many crop plants. Under saline conditions root growth has been found to be regulated by calcium and found to mitigate the adverse effect in barley (Cramer et al. 1990).

Pennisetum typhoides Stapf et Hubb. (Family: Poaceae), Pearl millet is one of the four most important cereals (rice, maize, sorghum and millets) grown in the tropics (The Syngenta Foundation for Sustainable Agriculture, 2002). Pearl millet is a crop of hot and dry climates, and can be grown in areas where rainfall is not sufficient (200-600 mm) for maize and sorghum. Global production exceeds 10 million tons a year

(National Research Council 1996). The food value of pearl millet is high. Trials in India have shown that pearl millet is nutritionally superior from human growth when compared to maize and rice. The protein content of pearl millet is higher than maize and has a relatively high vitamin A content.

The ameliorative effect of $CaCl_2$ during growth under NaCl stress requires detailed physiological studies in *P. typhoides* plants. This study aims at an understanding of the effect of salinity on the physiological parameters of *P. typhoides* under NaCl stress and also of the ameliorative effect of $CaCl_2$ on NaCl stressed plants.

MATERIALS AND METHODS

Plant material and growth: The seeds of *Pennisetum typhoides* were obtained from the M.S.Swaminathan Research Foundation, Kolli hills, Tamilnadu, India. The seeds were surface sterilized with 0.2% (w/w) $HgCl_2$ solution for 5min with frequent shaking and then thoroughly washed with deionised water. The seeds were sown in plastic pots (300mmdiameter) filled with 3 kg of soil mixture containing red soil, sand soil and farmyard manure at a 1:1:1 ratio. Ten seeds were sown per pot and all the pots were watered to field capacity with ground water up to 9 days after sowing. On 10 days after sowing the pots were irrigated with groundwater (control), 100mM NaCl, 100 mM NaCl with 5 mM $CaCl_2$ and 5 mM $CaCl_2$ solution. The plants were harvested randomly 15, 25 and 35 days after sowing and used for estimating the biochemical markers and antioxidant enzyme activities.

Biochemical analysis: The protein content was determined by the method of (Bradford 1976). Total free amino acids were extracted and estimated using the method of (Moore and Stein 1948). The proline content was estimated by the method of (Bates et al. 1973). The plant material was homogenized in 3%aqueous sulfosalicylic acid and the homogenate was centrifuged at 10000 g. Supernatant was used for estimation of proline content. The reaction mixture consisted of 2 ml acid ninhydrin and 2 ml of glacial acetic acid, which was boiled at 100 °C for 1 h. After termination of the reaction in an ice bath, the reaction mixture was extracted with 4 ml of toluene and absorbance was read at 520 nm. Soluble sugars were estimated from the samples by the method of (Dubois et al. 1956). Plant samples were homogenized in 80% ethanol (v/v). The homogenate was refluxed over a water bath and centrifuged. The residue was extracted twice more with 80% ethanol. The supernatant liquids were

combined and used for the estimation of soluble sugars.

Antioxidant enzyme extractions and assays:

Peroxidase activity (POX) was assayed by the method of (Kumar and Khan 1982). The assay mixture contained 2 ml of 0.1 M phosphate buffer (pH 6.8), 1 ml of 0.01 M pyrogallol, and 1ml of 0.005 M H₂O₂ and 0.5 ml of enzyme extract. The solution was incubated for 5 min at 25°C after which the reaction was terminated by adding 1 ml of 1.25 M H₂SO₄. The amount of purpurogallin formed was determined by measuring the absorbance at 420 nm against a blank prepared by adding the extract after the addition of 1.25 M H₂SO₄ at zero time. The activity was expressed in U mg⁻¹ protein. One U is defined as the change in the absorbance by 0.1 min⁻¹ mg⁻¹ protein. The assay of polyphenol oxidase (PPO) was carried out by the method of (Kumar and Khan 1982). The assay mixture contained 2 ml of 0.1 M phosphate buffer (pH 6.0), 1 ml of 0.1 M catechol and 0.5 ml of enzyme extract. This was incubated for 5 min at 25 °C, after which the reaction was stopped by adding 1ml of 1.25 M H₂SO₄. The absorbance of the quinone formed was read at 495 nm. To the blank 1.25 M H₂SO₄ was added of the zero time of the same assay mixture. PPO activity is expressed in U mg⁻¹ protein (U = Change in 0.1 absorbance min⁻¹ mg⁻¹ protein).

Catalase was measured according to the method of (Chandlee and Scandalios 1984) with small modifications. The assay mixture contained 2.6 ml of 50 mM potassium phosphate buffer (pH7.0), 0.4 ml of 15 mM H₂O₂ and 0.04ml of enzyme extract. The decomposition of H₂O₂ was followed by the decline in absorbance at 240 nm. The enzyme activity was expressed in U mg⁻¹ protein (U = 1 mM of H₂O₂ reduction min⁻¹ mg⁻¹ protein).

Statistical analysis: Each treatment (control, 100mM NaCl, 100 mM NaCl with 5 mM CaCl₂ and 5 mM CaCl₂ solution) was analysed with at least six replicates and a standard deviation (S.D.) was calculated and data are expressed in mean ± S.D. of six replicates.

RESULTS

Protein content: The NaCl lowered the protein content and CaCl₂ stress in root and shoot of the *Pennisetum typhoides* plants when compared with control plants. Addition of CaCl₂ with NaCl treatments increased the protein content when compared with NaCl stressed plants (Tab.1).

Amino acids and proline content: NaCl and CaCl₂ treatments increased the free amino acids and proline content in all parts of the plants as

compared to the control. CaCl₂ treatment to the NaCl stressed plants lowered the free amino acid and proline content in all parts when compared with NaCl treated plants (Tab.1). Among the organs the root showed a higher free amino acid content. Likewise the shoot showed increase in proline content.

Total soluble sugar content: NaCl and CaCl₂ treatments decreased the total sugar content in all organs when compared with control plants (Tab.1).When CaCl₂ was combined with NaCl treatments the plants showed increased the total sugar content when compared with NaCl treated plants. Among the organs, the roots showed a higher total sugar content, followed by a shoot in the treatment. NaCl treatment decreased the protein, sugar content and increased the free amino acid and proline content. CaCl₂ treatment improved all these solutes examined except for the free amino acids, which were decreased to the level of control by CaCl₂ treatments. Addition of calcium increased the protein and decreased the free amino acid in the salt-stressed plants (Tab.1).

Antioxidant enzyme contents: NaCl and CaCl₂ treated plants showed increased peroxidase activity as compared with the control. Combining CaCl₂ with the NaCl treatment decreased peroxidase activity as compared with NaCl treated plants (Fig. 1). Under NaCl and CaCl₂ stress, the activities of antioxidant enzymes like polyphenol oxidase (Fig. 2) and catalase (Fig. 3) decreased in all parts of the plants as compared to the control. When CaCl₂ was combined with NaCl, the treatment increased the polyphenol oxidase and catalase activity as compared with NaCl stressed plants. Among the organs, the roots showed a higher polyphenol oxidase and catalase activity followed by the shoot in all the treatments.

DISCUSSION

The NaCl lowered the protein content in sorghum (Azooz et al. 2004) and mungbean (Dhingra and Sharma 1993). Addition of CaCl₂ with NaCl treatments increased the protein content when compared with NaCl stressed plants. Similar results were observed in wheat (Abdlesamad 1993). Protein degradation in a saline environment has been attributed to the decrease in protein synthesis, accelerated proteolysis, decrease in the availability of amino acid and denaturation of enzymes involved in protein synthesis (Levitt 1980).

Free amino acids increased with an increase in NaCl salinity in *Catharanthus roseus* (Jaleel et al. 2007a) and CaCl₂ in wheat (Abdelasmad 1993). CaCl₂ treatment of the NaCl stressed plants lowered

the free amino acids and the proline content in all parts as compared with NaCl treated plants. Similar results were observed in the groundnut (Girija et al. 2002). Among the organs, the shoot showed a higher free amino acid and proline content followed by the root in all the treatments.


NaCl and CaCl₂ treatments decreased the total sugar content in *Cajanus cajan* (Gill and Sharma 1993). When CaCl₂ was combined with NaCl treatments, the plants showed an increase of the total sugar content as compared with NaCl treated plants. Among the organs, the roots showed a higher total sugar content followed by the shoot in the treatment. Addition of calcium may increase the α-amylase activity which leads to the degradation of starch into sugars. α-amylase requires Ca²⁺ for its activity (Bilderback 1973). Addition of calcium increased the protein and decreased the free amino acid in the Salt-stressed plants. So calcium treatment may reduce protein degradation or increase protein synthesis in salt-stressed plants.

Under salt stress, the antioxidant enzymes like polyphenol oxidase and catalase activities decreased in all parts of *Catharanthus roseus* (Jaleel et al. 2007a). When CaCl₂ was combined with NaCl, treatment increased the polyphenol oxidase and catalase activity as compared with NaCl stressed plants. Among the organs, the roots showed a higher polyphenol oxidase and catalase activity, followed by the shoot in all the treatments. The results obtained in this study were in accordance with those found in the roots of rice


seedlings (Lin and Kao 1999). The inhibition of POX activity by salinity may interfere with the regulation of auxin levels and also with cell-wall biosynthesis (Mittova et al. 2002). Low basal rate and decreased POX activity seem to indicate that this enzyme does not take a crucial part in defense mechanisms against oxidative stress or that, suffering POX for salt toxicity; cooperation is activated between different antioxidant enzymes for the establishment of a proper H₂O₂ homeostasis (Chaparzadeh et al. 2004). NaCl and CaCl₂ treated plants showed increased activity of peroxidase as compared with the control. Similar results were observed in spinach leaves (Ozturk and Demir 2003). Antioxidant enzymes play a significant role in plants to protect them against the damaging effect of reactive oxygen generated during salinity stress (Asada 1992). Reduction in catalase activity under salt stress may result in H₂O₂ accumulation and may be associated with its tolerant mechanism through signal transduction (Jaleel et al. 2007c). An increase in catalase activity was previously reported in triadimefon, a fungicide treated *Catharanthus roseus* plants (Jaleel et al. 2006).

Conclusion: NaCl was shown to have greater toxic effects than CaCl₂ on the growth and metabolism of *Pennisetum typhoides* plants. The addition of CaCl₂ to NaCl treatments had a varied effect on proteins, total sugars, free amino acids, proline and the activity of antioxidant enzymes in the growth of *Pennisetum typhoides* plants. In general Ca²⁺ was shown to have an ameliorative effect on NaCl stressed plant growth by modulating the overall metabolism.

Figures: 1. Effect of NaCl, CaCl₂ and their combinations on peroxidase (POX) activity in the (a) root and (b) shoot of *P.typhoides*. Values are given as mean ± SD of six samples in each group. DAS – days after sowing.



Figures: 2. Effect of NaCl, CaCl₂ and their combinations on polyphenol oxidase (PPO) activity in the (a) root and (b) shoot of *P.typhoides*. Values are given as mean ± SD of six samples in each group. DAS – days after sowing.



Figures: 3. Effect of NaCl, CaCl₂ and their combinations on catalase (CAT) activity in the (a) root and (b) shoot of *P.typhoides*. Values are given as mean ± SD of six samples in each group. DAS – days after sowing.




Table 1. Effect of NaCl, CaCl₂ and NaCl + CaCl₂ on proteins, free amino acids, proline, and total sugars content of *P.typhoides*. Values are given as mean \pm SD of six samples in each group. DAS – days after sowing.

Parameters	DAS	Root				Shoot			
		Control	100 mM NaCl	100 mM NaCl + 5 mM CaCl ₂	5mM CaCl ₂	Control	100 mM NaCl	100 mM NaCl + 5 mM CaCl ₂	5 mM CaCl ₂
Proteins mg g ⁻¹ DW	15	25.53 \pm 0.912	22.34 \pm 0.770	23.38 \pm 0.957	24.55 \pm 0.885	22.64 \pm 0.809	16.98 \pm 0.629	20.93 \pm 0.748	19.55 \pm 0.698
	25	33.71 \pm 1.162	29.72 \pm 1.061	31.58 \pm 1.169	32.17 \pm 1.149	31.57 \pm 1.089	25.83 \pm 0.891	29.74 \pm 1.062	28.71 \pm 1.063
	35	36.54 \pm 1.305	31.28 \pm 1.117	35.19 \pm 1.282	35.42 \pm 1.221	34.29 \pm 1.225	27.32 \pm 0.976	32.69 \pm 1.168	30.26 \pm 1.043
Free Amino Acid mg g ⁻¹ DW	15	6.284 \pm 0.022	7.153 \pm 0.247	6.165 \pm 0.220	6.639 \pm 0.237	3.119 \pm 0.104	3.673 \pm 0.127	2.826 \pm 0.097	3.109 \pm 0.109
	25	7.320 \pm 0.244	8.457 \pm 0.292	7.117 \pm 0.237	7.332 \pm 0.262	4.896 \pm 0.169	6.019 \pm 0.215	4.121 \pm 0.137	5.132 \pm 0.190
	35	8.165 \pm 0.292	9.219 \pm 0.307	8.054 \pm 0.259	8.274 \pm 0.285	6.452 \pm 0.215	7.286 \pm 0.260	5.176 \pm 0.185	6.982 \pm 0.249
Proline mg g ⁻¹ DW	15	0.419 \pm 0.015	0.518 \pm 0.019	0.474 \pm 0.016	0.498 \pm 0.018	0.468 \pm 0.017	0.593 \pm 0.019	0.513 \pm 0.018	0.554 \pm 0.019
	25	0.821 \pm 0.029	1.182 \pm 0.042	0.958 \pm 0.033	1.027 \pm 0.037	0.921 \pm 0.033	1.237 \pm 0.044	1.128 \pm 0.048	1.197 \pm 0.043
	35	0.941 \pm 0.033	1.326 \pm 0.046	1.119 \pm 0.039	1.274 \pm 0.046	1.079 \pm 0.037	1.496 \pm 0.053	1.227 \pm 0.042	1.329 \pm 0.048
Total Sugars mg g ⁻¹ DW	15	2.869 \pm 0.103	2.419 \pm 0.086	2.612 \pm 0.093	2.571 \pm 0.092	2.537 \pm 0.091	2.014 \pm 0.072	2.319 \pm 0.080	2.229 \pm 0.080
	25	4.782 \pm 0.165	4.123 \pm 0.142	4.327 \pm 0.156	4.216 \pm 0.150	4.487 \pm 0.160	3.758 \pm 0.134	4.297 \pm 0.154	4.107 \pm 0.147
	35	5.989 \pm 0.222	5.107 \pm 0.176	5.318 \pm 0.183	5.253 \pm 0.181	5.121 \pm 0.183	4.563 \pm 0.163	5.043 \pm 0.180	4.389 \pm 0.157

REFERENCES

1. Abdelsamad H.H. Counteraction of NaCl with CaCl₂ or KCl on pigment saccharide and mineral contents in wheat. Biol. Plant 1993; 35: 555–560.
2. Asada K. Ascorbate peroxidase – a hydrogen peroxide scavenging enzyme in plants. Physiol. Plant 1992; 55: 235–241.
3. Azooz M.M, Shaddad M.A, Abdel-Latef A.A. The accumulation and compartmentation of proline in relation to salt tolerance of three sorghum cultivars. Indian J. Plant Physiol 2004; 9: 1–8.
4. Bates L.S, Waldren R.P, Teare I.D. Rapid determination of free proline for water stress studies. Plant Soil 1973; 29: 205–207.
5. Bilderback D.E. A simple method to differentiate between alpha and beta-amylase. Plant Physiol 1973; 51: 594–595.
6. Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding. Ann. Biochem 72: 248–253.
7. Chandlee J.M, Scandalios J.G. Analysis of variants affecting the catalase development program in maize scutellum. Theor. Appl. Genet 1984; 69: 71–77.
8. Chaparzadeh N, Amico M.L, Nejad R.K, Izzo R, Izzo F.N. Antioxidative responses of Calendula officinalis under salinity conditions. Plant Physiol. Biochem 2004;42: 695–701.
9. Cramer G.R, Epstein E, Lauchli A. Effect of sodium, potassium and calcium on salt stressed barley. 1. Growth analysis. Physiol. Plant 1990; 80: 83–88.
10. Demiral T, Turkan I. Exogenous glycine betaine affects growth and proline accumulation and retards senescence in two rice cultivars under NaCl stress. Environ. Exp. Bot 2006; 56: 72–79.
11. Dhingra H.R, Sharma R.K. Biochemical and mineral composition of young healthy and shriveled mung bean seeds in response to salinity. Indian J. Plant Physiol 1993; 36: 115–117.
12. Dubois M, Gilles K.A, Hamilton J.K, Roberts P.A, Smith F. A calorimetric method for determination of sugars and related substances. Anal. Chem 1956; 28: 350–356.
13. Duncan R.R. Plant tolerance to acid soil constraints: genetic resources, breeding methodology and plant improvement. In Wilkinson, R.E. (ed.), Plant environment interactions, 2000; 1–38. Marcel Dekker, New York.
14. Flowers T.J, YEO A.R. Solute transport in plants. London: Blackie Academic and Professional 1992.
15. Gill KS, Sharma P.C. Mechanism of salt injury at seedling and vegetative growth stage in *Cajanus cajan* (L.) Millsp. Plant Physiol. Biochem 1993; 20: 49–52.
16. Girija C, Smith B.N, Swamy P.M. Interactive effects of sodium chloride and calcium chloride on the accumulation of proline and glycine betaine in peanut (*Arachis hypogaea* L.). Environ. Exp. Bot 2002; 47: 1–10.
17. Grattan S.R, Grieve C.M. Salinity – mineral nutrient relations in horticultural crops. Sci. Hort 1999; 78: 127–157.
18. Greenway H, Munns R. Mechanism of salt tolerance in non - halophytes. Ann. Rev. Plant Physiol 1980; 31: 149–190.
19. Hasegawa P.M, Bressan R.A, Zhu J.K, Bohnert H.J. Plant cellular and molecular responses to high salinity, Annu. Rev. Plant Physiol. Plant Mol. Biol 2000; 51: 463–499.

20. Hoshida H, Tanaka Y, Hibino T, Hayashi Y, Tanaka A, Takabe T. Enhanced tolerance to salt stress in transgenic rice that over expresses chloroplast glutamine synthetase. *Plant Mol. Biol* 2000; 43: 103–111.
21. Jaleel C.A, Gopi R, Alagu Lakshmanan G.M, Panneerselvam R. Triadimefon induced changes in the antioxidant metabolism and ajmalicine production in *Catharanthus roseus* (L.) G. Don. *Plant Sci* 2006; 171: 271–276.
22. Jaleel C.A, Gopi R, Manivannan P, Panneerselvam R. Antioxidative potentials as a protective mechanism in *Catharanthus roseus* (L.) G. Don. *Plants under salinity stress*. *Tur. J. Bot* 2007b; 31: 245–251.
23. Jaleel C.A, Gopi R, Manivannan P, Panneerselvam R. Responses of antioxidant defense system of *Catharanthus roseus* (L.) G. Don. To paclobutrazol treatment under salinity. *Acta Physiol. Plant* 2007c; 29: 205–209.
24. Jaleel C.A, Gopi R, Sankar B, Manivannan P, Kishorekumar A, Sridharan R, Panneerselvam R. Studies on germination, seedling vigour, lipid peroxidation and proline metabolism in *Catharanthus roseus* seedlings under salt stress. *S. Afr. J.Bot* 2007a;73: 190–195.
25. Kumar K.B, Khan P.A. Peroxidase and polyphenol oxidase in excised ragi *Eleusine coracana* (V. PR 202) leaves during senescence. *Indian J. Exp. Bot* 1982; 20: 412–416.
26. Lawlor D.W. Limitation to photosynthesis in water stressed leaves: Stomata Vs metabolism and role of ATP. *Ann. Bot* 2002; 89: 871–885.
27. Levitt J. Responses of plants to environmental stresses. Academic Press, New York 1980.
28. Lin C.C, Kao C.H. NaCl induced changes in ionically bound peroxidases activity in roots of rice seedlings. *Plant Soil* 1999; 216: 147–153.
29. Mittler R. Oxidative stress, antioxidants and stress tolerance, *Trends Plant Sci* 2002; 7:405–410.
30. Mittova V, Guy M, Tal M, Volokita M. Response of the cultivated tomato and its wild salt-tolerant relative *Lycopersicon pennellii* to salt-dependent oxidative stress; increased activities of antioxidant enzymes in root plastids. *Free Radical Res* 2002; 36: 195–202.
31. Moore S, Stein W.H. Photometric method for use in the chromatography of amino acids. *J. Biol. Chem* 1948; 175: 367–388.
32. Munns R. Genes and salt tolerance: bringing them together. *New Phytol* 2005; 167: 645–663.
33. Munns R. Physiological processes limiting plant growth in saline soils: some dogmas and hypotheses, *Plant Cell Environ* 1993; 16:15–24.
34. National Research Council, Lost crops of Africa. Volume I: Grains. National Academy Press. Washington DC. 1996.
35. Nemoto Y, Sasakuma T. Differential stress responses of early salt-stress responding genes in common wheat. *Phytochemistry* 2002; 61: 129–133.
36. Oudhia P, Tripathi R.S. Medicinal weed flora of brinjal (*Solanum melongena* L.) Hisar region. *Crop Res* 2002; 20: 482–488.
37. Ozturk L, Demir Y. Effects of putrescine and ethephon on some oxidative stress enzyme activities and proline content in salt stressed spinach leaves. *Plant Growth Regul* 2003; 40: 89–95.
38. Reddy V.S, Reddy A.S.N. Proteomics of calcium-signaling components in plants. *Phytochemistry* 2004; 65: 1745–1776.
39. Rengel Z. The role of calcium in salt toxicity. *Plant Cell Environ* 1992; 15: 625–632.
40. Sairam R.K, Srivastava G.C. Changes in antioxidant activity in sub-cellular fractions of tolerant and susceptible wheat genotypes in response to long term salt stress. *Plant Sci* 2002; 162: 897– 904.
41. Shakila A, Rajeswari R. Effect of GA₃, certain chemicals and media on germination, growth and yield of *Phyllanthus niruri*. *Plant Arch* 2006; 6: 121–125.
42. Sohan D, Jason R, Zajcek J. Plant-water relations of NaCl and calcium-treated sunflower plants. *Environ. Exp. Bot* 1999; 42:105–111.
43. Teixeira J, Pereira S. High salinity and drought act on an organ dependent manner on potato glutamine synthetase expression and accumulation. *Environ. Exp. Bot* 2007; 60: 121–126.
44. The Syngenta Foundation for Sustainable Agriculture, Millet: Pearl Millet. [http://syngentafoundation.org/what is pearl millet.htm](http://syngentafoundation.org/what_is_pearl_millet.htm). 2002.
45. Yamaguchi T, Blumwald E. Developing salt- tolerant crop plants: challenges and opportunities. *Trends Plant Sci* 2005; 10: 615–620.