



Assessment of knowledge, awareness and reporting of pharmacovigilance among the Nursing professionals

Duarairajan P^{1*}, Sujitha M² and Maheshwari Jaikumar³

¹Professor & HOD, ²Tutor, Department of Pharmacology, Sri Muthukumar Medical College Hospital & Research Institute, Chikkararyapuram (Near Mangadu), Chennai – 600069 [Affiliations to the Tamil Nadu Dr. M.G.R Medical University, Chennai, India].

³Professor & Principal, Meenakshi College of Nursing, MAHER (Declared as Deemed to be University U/S 3 of UGC Act 1956), Chikkarayapuram (Near Mangadu), Chennai- 600069

Received: 19-04-2018 / Revised Accepted: 17-05-2018 / Published: 01-06-2018


ABSTRACT

There is a considerable increase in awareness about the issues related to drug safety among the Nursing professionals, healthcare institutions and the public. Our aim of this present study was to assess the knowledge; awareness and reporting of adverse drug reactions among the Nursing professionals. Methodology: A questionnaire model was prepared, focusing on the awareness, knowledge of various adverse drugs reactions and the algorithm of reporting systems of pharmacovigilance programme of India. A total of 100 Nursing professionals were included in this study. The questionnaire was distributed among the Nursing Faculties working in various specialties. Completed questionnaire were collected and analysed statistically. Data collected was analysed and was reported as percentage of responders and non-responders. Total responders were 61% and 39% were non-responders. This indicates that the knowledge of ADR reporting was not up to the mark which emphasizes need to implement, appropriate strategies to strengthen the awareness of pharmacovigilance practices and the importance of ADR reporting in this hospital.

Keywords: Pharmacovigilance Programme of India, Awareness, Knowledge, Reporting Nursing professionals

Address for Correspondence: Dr. P. Durai Rajan M.D., Professor & HOD, Department of Pharmacology, Sri Muthukumar Medical College & Research Institute, Chennai, Tamilnadu, India; Email: drpdrsrc@gmail.com

How to Cite this Article: Duarairajan P, Sujitha M and Maheshwari Jaikumar. Assessment of knowledge, awareness and reporting of pharmacovigilance among the Nursing professionals. World J Pharm Sci 2018; 6(6): 107-111.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License, which allows adapt, share and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms. 

INTRODUCTION

World Health Organization defines pharmacovigilance as “the science and activities relating to the detection, evaluation, understanding, reporting and prevention of adverse drug effects or any other drug related problems. The term pharmacovigilance has evolved to recognize the importance for monitoring and improving the safe use of medicines. The burden of adverse drug reactions (ADRs) in the global scenario is highly accountable for considerable morbidity, mortality, and financial commitment to the patients [1,2]. In England, 0.9% of the total hospital admissions were due to ADRs during the year 1999–2008. ADRs are common in Australian healthcare system also and they contribute to 1% of hospital admissions. In the United States of America, ADRs contribute 3.4%–7% of hospital admissions. The percentage of hospital admissions due to ADRs in certain countries is 10% or more [3]. India, with a current population of 1.27 billion, is the fourth largest producers of pharmaceuticals in the world with more than 6000 licensed manufacturers and over 60,000 branded formulations in the market. Studies revealed that ADRs are leading to hospitalization and constitute a significant economic burden on patients in India. A study showed that hospital admissions due to ADRs accounted for 0.7% of total admissions and deaths due to ADRs accounted for 1.8% of total admissions in a territory referral center in South India [4,5].

Promoting safe use of medicines is a priority of Indian Pharmacopoeia Commission that functions as the National Coordination Center (NCC) for Pharmacovigilance Programme of India (PvPI). One hundred and seventy-nine adverse drug reactions (ADRs) monitoring centers currently report ADRs to NCC. Current India contribution to global safety database reaches 3% and the completeness score is 0.93 out of 1. NCC is taking several measures to enhance patient safety including capacity building for monitoring, surveillance, collaboration with national health programs and other organizations to increase ADR reporting and to ensure that PvPI is a vital knowledge database for Indian regulators. The z Central Drugs Standard Control Organization has notified important safety label changes on drugs such as carbamazepine and piperacillin + tazobactam in the year 2015, other drugs are under monitoring for regulatory interventions [6].

In developing countries such as India, under-reporting of ADR remains a serious concern. In India the Information and Technology (IT) is becoming a great facilitator for promoting public health. India is cementing its place at IT sector

through mobile connections to reach every individual in a population of 1.27 billion, where approximately 77.58% population is already using mobile phones. Hence, it is more rationale to introduce the concept of PvPI to stakeholders through mobile phones. NCC- PvPI in technical collaboration with NSCB Medical College, Jabalpur, developed a mobile application for the healthcare professionals to promote easy and instant reporting of ADR. This facility was launched by Secretary Health, MoHFW, Government of India, on May 22, 2015 [7,8]. To improve the active participation of patients, healthcare professionals, and the pharmaceutical industrial reporting of suspected ADRs to the PvPI, NCC recently launched a helpline number (toll free), i.e., 1800 180 3024 facility for reporting adverse events. This facility was dedicated to the nation on October 11, 2013. This may be one of the innovative methods to create awareness in each and every corner of the country for the pharmacovigilance activity. This facility will be useful for the healthcare professionals who are working in tertiary healthcare system and also who are working in other health care functionary for easy reporting of ADRs. Adverse events-related information received at NCC and the same will be communicated to the nearest AMCs for analysis and validation of the reports. Since sending timely feedback or acknowledgement will build up public confidence and this facility has been upgraded by sending short message service feedback/acknowledgment to the ADRs reporters [9].

METHODOLOGY

About 100 Nursing professionals were included in the study. Questionnaire developed by PvPI to assess the knowledge, awareness and reporting of pharmacovigilance, awareness of Pharmacovigilance programme of India. The Questionnaire of PvPI were distributed to all the participants. The questionnaire model was clearly explained to each and every participant. Sufficient time was given to fill up the feedback forms. Completed feedback forms were collected from the participants. The feedback forms were divided into responders and non-responders. The results were analyzed on the basis of number who has responded correctly by marking yes / no and the same was reported as percentage for statistical significance.

RESULTS

Statistical analysis of the study is shown in table 1. Which showed the responses to the Question number 1 was 100%. The response to question number 2,3,4,5,6,8,9 and 10 were around 60% to 95% which was satisfactory. Whereas the response

to questions 7,11 and 12 were below 50%, and the same is shown clearly in fig:1, which was not up to the mark.

DISCUSSION

The positive response to question number one regarding adverse drug reactions and their personal experience at any time following any medicinal use was satisfactory whereas the response for knowledge, relating to existence of pharmacovigilance programme of India (PvPI), for safe use of medicines and if at all any adverse drug reactions occurs, when, to whom, where to report and what are the step wise procedures involved in reporting were only below 40% as shown in table 2. This study is not in accordance with one previous study, where only 2.7% were aware, and ignorance about ADRs reporting were 71.4%, address to report 68% unaware to report 52.2% and needless to report were 44.4%[10]. The awareness and knowledge of various pharmacovigilance programme of India monitoring centers in their region, the specifically designed formats for reporting ADRs and the availability of toll free helpline number (**1800-180-3024**) to report any suspected ADRs after the use of medicines and medicine safety promotional materials, were below 20% only. All participants were very eager to participate PvPI [11].

The burden of adverse drug reactions in medicine safety initiatives of (ADRs) in the global scenario is high and accounts for considerable morbidity, mortality, and extra – cost to the patients [12, 13]. In England, 0.9% for the total hospital admissions was due to ADRs during the year 1999 – 2008. ADRs are common in Australian healthcare system also and they contribute to 1% of hospital admissions. In the Untied States of America, ADRs in contribute 3.4% - 7% of hospital admissions. The percentage of hospital admissions due to ADRs in certain countries is 10 % or more. India, with a current population of 1.27 billion, is the fourth largest producers of pharmaceuticals in the world with more than 6000 licensed manufactures and over 60,000 branded formulations in the market. Studies revealed that ADRs are leading to hospitalization and constitute a significant economic burden on patients in India. Various studies have showed that hospital admissions due to ADRs accounted for 0.7% of total admissions and deaths due to ADRs accounted for accounted for 1.8% of total admissions in a territory referral center in South India [14].

Therefore, medicines safety monitoring is an essential element of healthcare and for high quality medical care. Since safety monitoring of medicines as an integral part of clinical practice, the Ministry of Health of Health and Family Welfare (MoHFW), Government of India launched the nationwide Pharmacovigilance Programme of India (PvPI) in the year 2010 to inspire confidence and trust among patients and healthcare professionals with respect to medicines safety. Indian Pharmacopoeia Commission (IPC) under the MoHFW has been functioning as the National Coordination centre (NCC) for PvPI since April 2011. There has been rapid progress in reporting of ADRs by the healthcare professionals in the past 5 years [15].

Periodic safety update reports (PSURs) are a tool to monitor the safety of ongoing medicines in the market. In India, marketing authorization holders (MAHs) are required to prepare PSURs and to submit them to CDSCO twice in a year for 2 years and annually for another 2 years [16]. Since PSURs are not directly linked with PvPI, NCC has taken the initiatives in collaboration with CDSCO to utilize the data for PvPI. The first interactive session of “Review of Periodic Safety Update Report/Post Marketing Surveillance Date and

Pharmacovigilance planning of Marketed Produced” was held on December 18 and 19, 2013 at New Delhi. Representatives from MAHs, CDSCO, and NCC-PvPI had participated and discussed the issues and roadmap for better coordination and participation of MAHs in PvPI [17].

CONCLUSION

The present pharmacovigilance study demonstrates the lack of awareness and knowledge among the Nursing professionals. Our findings provide information to healthcare policy makers and healthcare authorities in the future for the evaluation and reinforcement of plans to improve the pharmacovigilance awareness.

ACKNOWLEDGMENT

The authors would like to thank the Management and the Dean, the Vice Principal Sri Muthukumaran Medical College Hospital and Research Institute for their help in conducting and publication of the study. We thank the principal, Meenakshi College of Nursing to get feedback from the faculties. We also thank our department staffs for their continued support.

TABLE: 1 SHOWING POSITIVE AND NEGATIVE RESPONDERS


Questions	Number of participants	Positive response	Negative response
1	100	100	0
2	100	93	7
3	100	84	16
4	100	52	48
5	100	62	38
6	100	52	48
7	100	25	75
8	100	75	25
9	100	61	39
10	100	64	36
11	100	33	67
12	100	28	72

Table: 1 shows Qn:1,2,3 Positive Response 100,93,84 and Negative Response 0,7,16. Qn:4,5,6 shows Positive Response 52,62,52 and Negative Response 48,38,48. Qn:7,8,9 shows Positive Response 25,75,61 and Negative Response 75,25,39. Qn:10,11,12 shows Positive Response 64,33,28 and Negative Response 36,67,72.

Table - 2: SHOWING THE PERCENTAGE OF POSITIVE&NEGATIVE RESPONSE

Questions No	Positive response	Percentage (%)	Negative response	Percentage (%)
1	100	100	0	0
2	93	93	7	7
3	84	84	16	16
4	52	52	48	48
5	62	62	38	38
6	52	52	48	48
7	25	25	75	75
8	75	75	25	25
9	61	61	39	39
10	64	64	36	36
11	33	33	67	67
12	28	28	72	72

Table: 2 shows Qn:1,2,3 shows percentage of positive response 100,93,84 Negative response 0,7,16. Qn:4,5,6 shows percentage of positive response 52,62,52 Negative response 48,38,48. Qn:7,8,9 shows percentage of positive response 25,75,61 Negative response 75,25,39. Qn:10,11,12 shows percentage of positive response 64,33,28 Negative response 36,67,72.



REFERENCES

1. The World Health Organization, Safety of medicines: A guide to detecting and reporting adverse drug reactions 2002a; Geneva.
2. Ernst F.R., et al. Drug-related morbidity and mortality: updating the cost-of-illness model. *J. Am. Pharm. Assoc* 2001; 41:192–199.
3. Lazarou J., et al. The incidence of adverse drug reactions in hospitalized patients—a meta-analysis of prospective studies. *JAMA* 1998; 279:1200–1205.
4. Classen D.C., et al. Adverse drug events in hospitalized patient—excess length of stay, extra cost and attributable mortality. *JAMA* 1997; 277:301–306.
5. Ahmed M.el-B. et al. Drug-associated admissions to a district hospital in Saudi Arabia. *J. Clin. Pharm. Ther.* 1997; 22:61–66.
6. The World Health Organization, The importance of pharmacovigilance: Safety Monitoring of Medicinal Products 2002b; Geneva.
7. Belton K.J. et al. Attitude survey of adverse drug-reaction reporting by health care professionals across the European Union. The European Pharmacovigilance Research Group. *Eur. J. Clin. Pharmacol.* 1997; 52:423–427.
8. Herdeiro M.T., et al. Physicians' attitudes and adverse drug reaction reporting: a case-control study in Portugal. *Drug Saf* 2005; 28:825–833.
9. Lee A., Thomas S.H.L. Adverse drug reactions. In: Walker R., Edward C., editors. *Clinical Pharmacy and Therapeutics*. 3rd ed. Churchill Livingstone, 2003; 33–46.
10. Li Q., et al. Awareness and attitudes of healthcare professionals in Wuhan, China to the reporting of adverse drug reactions. *Chin. Med. J. (Engl.)* 2004; 117:856–861.
11. Cosentino M., et al. Attitudes to adverse drug reaction reporting by medical practitioners in a Northern Italian district. *Pharmacol. Res.* 1997; 35:85–88.
12. Graille V., et al. Drug vigilance: opinion survey among residents of a university hospital, *Therapie* 1994; 49:451–454.
13. Pirmohamed M., et al. Adverse drug reactions as cause of admission to hospital: prospective analysis of 18820 patients. *BMJ* 2004; 329:15–19.
14. Jick H. et al. Adverse drug reactions: the magnitude of the problem. *J. Allergy Clin. Immunol.* 1984; 74:555–557.
15. Al-Hazmi N.N., et al. A study of community pharmacists' and contributions to adverse drug reactions (ADRs) reporting systems in the Makkah, Kingdom of Saudi Arabia (KSA) *J. Clin. Trials* 2013; 3:127–132.
16. WHO Programme Members-Uppsala Monitoring Centre, Sweden: www.who-umc.org
17. Feely J., et al. Stimulating the reporting of an adverse drug reaction by using a fee. *Br. Med. J.* 1990; 300:22–23.