



---

## ***Buchanania Lanza*: a species of enormous potentials**

Mahtab Zakira Siddiqui<sup>1\*</sup>, Arnab Roy Chowdhury<sup>1</sup>, Niranjana Prasad<sup>1</sup> and Moni Thomas<sup>2</sup>

<sup>1</sup>Processing and Product Development Division, Indian Institute of Natural Resins & Gums, Namkum, Ranchi – 834 010 (Jharkhand), India

<sup>2</sup>Directorate of Research Services, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur – 482 004 (M.P.), India

---

Received: 17-02-2014 / Revised: 24-02-2014 / Accepted: 25-03-2014

---

### **ABSTRACT**

*Buchanania lanzan* Spreng, commonly known as char, achar and chironji, belongs to family Anacardiaceae. It was first described by Francis Hamilton in 1798. The tree is natural wild growth in the tropical deciduous forests of Northern, Western and Central India, mostly in the States of Chhattisgarh, Jharkhand, Madhya Pradesh and in Varanasi and Mirzapur districts of Uttar Pradesh. Besides India, the plant is also found in other tropical Asian countries, Australia and Pacific islands. Tree can be identified by dark grey crocodile bark with red blaze and is a good species for afforestation in bare hill slopes. Traditional indigenous knowledge reveals the immense value of almost all parts of the plant i.e. roots, leaves, fruits, seeds and gum for various medicinal uses. *Buchanania lanzan*, being a vulnerable medicinal plant, is included in the Red Data Book published by International Union for Conservation of Nature and Natural Resources (IUCN). This species has high socio-economic value providing livelihood to tribal population of the area and has high potential as commercial horticulture species. Unfortunately, due to over-exploitation and indiscriminate harvesting (lopping and cutting), leading to very severe threat to its extinction, which call for an urgent conservation effort at all levels. The production of this economically important forest tree species is further threatened by insect pests also. In this background, there is compelling need for developing a suitable technology facilitating easy multiplication, regeneration and conservation of the species, simultaneously imparting and disseminating proper knowledge and education to the tribal population.

**Key words:** *Buchanania lanzan*, piyar, chironji, char, potential uses.

---

### **INTRODUCTION**

*Buchanania lanzan* Spreng, commonly known as char, achar and chironji, belongs to family Anacardiaceae. It was first described by Francis Hamilton in 1798. It is endemic in the dry deciduous tropical forests of India and is an ever-green moderate-sized tree, with straight, cylindrical trunk, upto 10-15 m height and tomentose branches. Its bark is rough, dark grey or black, fissured into prominent squares, 1.25 to 1.75 cm thick, and is reddish inside. Flowering starts in the month of November and its leaves are coriaceous, broadly oblong with a rounded base. It bears fruits, each containing a single seed known as 'chironji' and is quite popular as an edible nut. It avoids waterlogged areas, but occurs on yellow sandy loam soils. Tree can be easily identified by its dark

grey crocodile bark with red blaze and is a good species for afforestation in bare hill slopes. *Buchanania lanzan*, being a vulnerable medicinal plant, is included in the Red Data Book published by International Union for Conservation of Nature and Natural Resources (IUCN) [1,2]. Regionally, it is known by different names. The botanical origin and vernacular names of *Buchanania lanzan* are given in the Table 1.

Chironji originated in the Indian sub-continent. The tree is found as natural wild growth in the tropical deciduous forests of Northern, Western and Central India, mostly in the States of Chhattisgarh, Jharkhand, Madhya Pradesh and in Varanasi and Mirzapur districts of Uttar Pradesh [3]. Besides India, the plant is found distributed in other tropical Asian countries, Australia and Pacific islands also.

---

\*Corresponding Author Address: Dr. (Ms) Mahtab Z. Siddiqui, Senior Scientist, Processing and Product Development Division, Indian Institute of Natural Resins & Gums, Namkum, Ranchi – 834010 (Jharkhand), India: E-mail: [mzs\\_2009@rediffmail.com](mailto:mzs_2009@rediffmail.com)

The species has high socio-economic value providing livelihood to tribal population of these areas and has high potential as commercial horticulture species. It is a common associate of sal (*Shorea robusta*), teak (*Tectona grandis*), dhok/kaldhi (*Anogeissus pendula*) and salai (*Boswellia serrata*). About seven species of *Buchanania* have been reported in India, out of which *Buchanania lanzan* and *Buchanania axillaries* (Syn. *Angustifolia*) produce edible fruits. *Buchanania lanceolata*, an endangered species, is found in the evergreen forests of Kerala while *Buchanania platyneura* is found in Andaman. Other species of the genus are *Buchanania lucida*, *Buchanania glabra* and *Buchanania acuminate*.

#### **Traditional indigenous uses**

Traditional indigenous knowledge reveals the immense value of almost all parts of the plant like roots, leaves, fruits, seeds and gum for various medicinal uses. The roots are acrid, astringent, cooling, depurative and constipating and are useful in treatment of diarrhea. Extract of the root is also used as an expectorant and for curing biliousness and blood diseases. The leaf juice is used as expectorant, aphrodisiac, purgative, blood purifier, thirst-quencher and cures digestive disorders. It contains 2.64% tannins (0.35% gallo-tannins), triterpenoids, saponins, flavonoids and reducing sugars. Powdered or crushed leaves are applied to wounds. The chironji seeds/kernels are nutritional, palatable and used as a substitute for almonds in confectionery. The seeds possess 3.0% moisture and are rich in lipid/fat (59.0%), protein (19.0–21.6%), starch/carbohydrate (12.1%), fibre (3.8%), minerals like calcium (279.0 mg), phosphorus (528.0 mg), iron (8.5 mg) and vitamins like thiamine (0.69 mg), ascorbic acid/vitamin C (5.0 mg), riboflavin (0.53 mg), niacin (1.50 mg) and also contain 34-47 % fatty oil which is used as a substitute for olive and almond oils. The calorific value of kernel is 650 k-cal/100g. Kernel lipids comprised mainly of neutral lipids (90.4%), consist mostly of triacylglycerol (82.2%), free fatty acids (7.8 %) and small amount of diacylglycerols, monoacylglycerols and sterols. Ointment made from its kernels is used to relieve itch and prickly heat. The oil extracted from kernels is known as 'char' and used for curing skin diseases and removing spots/blemishes from the face. The oil is also applied externally on glandular swellings of the neck singly, as also in combination with other herbal oils. Chironji seeds are tolerant to desiccation and chilling and show 95-100% survival upto 90 days at all storage temperatures with gradual loss in germination after 280 days of storing [4-7]. The seeds are used for preparing a traditional sweet dish known as 'chirongi ki burfi'. Average annual seed collection is 300 to 1200

quintals in Madhya Pradesh alone. It is an income-generating produce for the forest dependent communities. On an average, 40–50 kg fresh fruits are produced per tree, which come down to 8–10 kg on drying, yielding 1.0–1.5 kg of finished produce per tree [8]. The fruits are laxative and are used to relieve thirst, body-burning, fever, cough and asthma. Bark yields tannin (upto 13%) which is used in the tannin industries. The tribals of Southern Bihar blend the powder of the stem bark and *Syzygium cumini* (Myrtaceae) together and the same is given to treat infantile diarrhea. The gum oozed from the cut-bark is soluble in water and used internally for treatment of intercostals pain and diarrhea. The gum is mixed with goat's milk for effective and curative results in intercostal pains, being analgesic. Some tribal communities of Andhra Pradesh consume a blend of the gum dissolved in cow's milk for treating rheumatic pains. It is believed that the production and collection of about 175 metric tons of char gum is from Mandla, Dindori, Umariya, Shahdol, Katni and Chhindwara districts of Madhya Pradesh (India). The gum is generally used for adulteration of guggul (*Commiphora wightii*) by adding some perfume/scent. In the tobacco industry it is used for the refinement. The superior quality gum is used in soft drinks and edibles for coloring.

The timber of chironji is slightly resistant to termite and is utilized for making furniture, boxes and crates, desks, fine furniture, match boxes, moulding, packing cases, stools, tables and agricultural implements. The tribal communities of Sonbhadra District, one of the most backward districts of Uttar Pradesh and a part of the Vindhyan zone, earn money by collecting gum and lac by rearing kusumi strain of lac on the chironji trees. Thus, *Buchanania lanzan* is a socio-economically important underutilized life-support and tropical medicinal species for the tribal populace of North, West and Central India. But unfortunately due to over-exploitation and indiscriminate harvesting (lopping and cutting) considerable reduction in the population of *Buchanania lanzan* has been recorded in the recent past, leading to severe threat of its extinction, which needs urgent conservation efforts [9-11]. The production of this economically important forest tree species is threatened by insect pests also. Of them, stem borer *Plocaederus obesus* is the most harmful insect pest in Central India [12-15].

#### **RESEARCH LITERATURE REVIEW**

**Seed/kernel:** Sengupta and Roychoudhury in 1977 [16] reported the fatty acid composition of *Buchanania lanzan* seed oil by urea complex formation, using gas liquid chromatography (GLC).

Their study revealed that *Buchanania lanzan* seed oil contained myristic (0.6%), palmitic (33.4%), stearic (6.3%), oleic (53.7%) and linoleic (6.0%) acids and it might be a promising commercial source of palmitic and oleic acids. Determination of lipid composition of chironji kernel revealed the presence of neutral lipids (90.4%), glycolipids (3.4%) and phospholipids (6.2%). Neutral lipids consisted mostly of triacylglycerols (82.2%), free fatty acids (7.8%) and small amount of diacylglycerols, monoacylglycerols and sterols as also three glycolipids and six phospholipids [17].

The *in vivo* anti-inflammatory activity of methanolic extract of *Buchanania lanzan* kernel using carrageenan-induced paw edema as an acute model and the formaldehyde induced arthritis as a chronic model, has been reported by Warokar *et al.* in 2010 and they found that the 200 mg/kg body weight of extract significantly decreased paw volume. They also studied the *in vitro* antioxidant activity of the methanolic extract by 1,1-diphenyl-2-picryl hydrazyl (DPPH) method and found that 125 µg/ml of extract exhibited 83.11 and 88.34% scavenging DPPH and % reducing power respectively as compared with the standard antioxidant ascorbic acid (99.58%, 98.92%) [18]. Feeding of ethanolic extracts of dry fruits of *Prunus amygdalus* (almond) and *Buchanania lanzan* significantly stimulated both cell mediated immunity (CMI) and humoral immunity in BALB/c mice as evidenced by the enhancement of macrophage migration index (MMI), haemagglutinating antibody (HA) titres, and plaque forming cell (PFC) counts suggesting that these could be given to mothers after child birth or to invalids for enhancing their immunity [19]. The antioxidant activity and total phenolic content of methanolic extract of different dry fruits viz. almond, walnut, cashew nut, raisin and chironji by several chemical and biochemical assays including reducing power, lipid peroxidation damage in biomembranes, determination of antioxidant enzymes activity have been studied by Mishra *et al.* They found that methanolic extract of walnut showed higher value of antioxidant activity based on lipid peroxidation assay.

The higher phenolic content was found in walnut followed by almond, cashew nut, chironji and least in raisin. Walnut revealed the best antioxidant property, presenting lower EC<sub>50</sub> values in all assays except in antioxidant enzymatic activity [20]. Isolation of a novel and potent bio-material from the seeds of *Buchanania lanzan* and evaluation of its bio film forming ability, by formulating various ophthalmic films using PEG-400 as plasticizer, and the ultimate drug release studies from the formulated ophthalmic films exhibited a promising

stability, swelling index, folding endurance and sustainability for a period of 8 hrs., establishing that the isolated bio-material could act as a novel film former for formulating various ophthalmic films [21]. Tyagi and Madhav [22] reported the formulation of three different zidovudine micro emulsions, using the bio-emulsifier isolated from the seeds of *Buchanania lanzan* and the formulated micro-emulsions exhibited promising transparency, stability, uniform globule size and shape and surface tension for a period of 24 hrs. during drug release studies.

**Fruit:** Rai reported the methods of propagation and estimation of fruit yield of *Buchanania lanzan* Spreng. in 1982 [23]. The wound healing effect of alcoholic extract of fruits of *Buchanania lanzan* and its effect in dexamethasone suppressed wound healing was studied in Albino rats using three types of wound models i.e. incision, excision and dead space wounds and it was reported that the presence of flavonoids in *Buchanania lanzan* enhanced pro-healing activity [24].

**Leaves:** The leaves of *Buchanania lanzan* are reported to have immense medicinal value. Preliminary phytochemical investigation on leaves of *Buchanania lanzan* using TLC (Thin Layer Chromatography) and HPTLC (High Performance Thin Layer Chromatography) techniques revealed the presence of flavonoids, tannins, glycosides, phenols, steroid, saponin and myricetin 3'-rhamnoside-3-galactoside [25-28]. Mitra and Mehrotra [29,30] carried out pharmacognostical studies of leaf and bark of *Buchanania lanzan* Spreng. in 1981 and its fruits and seeds in 1982 [31]. Verma *et al.* [32] have reported that the methanolic extract of leaves of *Buchanania lanzan* Spreng. was more effective against gram negative bacteria (*Vibrio cholerae* and *Klebsiella pneumoniae*) when tested by paper disc diffusion method. Similarly, Manjunath and Mithun have also found that the methanolic extract of leaves of *Buchanania lanzan* was more significantly active against *Escherichia coli*, *Salmonella typhi*, *Staphylococcus aureus*, *Bacillus subtilis* in comparison to extracts of *Buchanania lanzan* leaves in petroleum ether, chloroform and water [33]. Adaptogenic (anti-stress) activity of methanolic extract of leaves of *Buchanania lanzan* using the swim endurance model in all groups under normal and stressed conditions was carried out by Mehta *et al.* The methanolic extract exhibited significant anti-stress activity in a dose-dependent manner [34]. Evaluation of antidiabetic, antihyperlipidemic and antioxidant activity of the methanol extract of leaves of *Buchanania lanzan* in streptozotocin-induced types I and II diabetic rats have been reported by Sushma *et al.* in 2013 [35].

**Bark:** Chaudhary *et al.* in 2001 studied the effect of aqueous extract of the bark of *Buchanania lanzan* Linn. on behaviour and chromatophores of a fresh water fish, Labeorohita and observed that in higher doses 450 ppm and 550 ppm fish could survive only for 76 hr. and 4.30 hr. respectively due to the toxicity of plant. They have reported that the doses ranging from 150 ppm to 350 ppm were psychoactive in nature [36]. Curative properties of methanolic extract of *Buchanania lanzan* bark as evaluated by its antioxidant, anti-inflammatory and DNA protective properties revealed that the methanolic extract could scavenge 2,2'-azino-bis (3-ethyl benzothiazoline-6-sulphonic acid) (ABTS) radicals with IC<sub>50</sub> of 0.25 mg/ml. The extract also significantly inhibited 15-lipoxygenase (15-LOX) and human cyclooxygenase (COX-2) in a dose-dependent manner, besides abolishing elastase activity. Further, the methanolic extract could prevent damage to DNA from the hydroxyl radicals produced during Fenton reaction [37]. Presence of chemo-protective phytoconstituents in the ethanolic extract of bark of *Buchanania lanzan* Spreng. offering protection against cyclophosphamide induced genotoxicity and oxidative stress in mice have been reported by Jain and Jain in 2012 [38]. The *in vitro* screening of methanolic extracts of the leaf and bark of *Buchanania lanzan* was carried out using DPPH, ABTS and H<sub>2</sub>O<sub>2</sub> (Hydrogen peroxide) radicals. The bark and leaf extracts exhibited potent inhibition against ABTS radical generation. The extracts showed good activity against DPPH radical, but in H<sub>2</sub>O<sub>2</sub> method leaf extract failed to show inhibition even at highest test concentration, whereas the bark extract, inhibited H<sub>2</sub>O<sub>2</sub> moderately [39].

**Gum:** Bothara and Singh have studied the characterization of thermal property of natural gums obtained from the seeds of *Diospyros melonoxylon* Roxb., *Buchanania lanzan* Spreng. and *Manilkara zapota* Linn. using differential scanning calorimetry (DSC), differential thermal analysis (DTA) and thermogravimetric analysis (TGA) under nitrogen atmosphere. They have reported that these gums are thermally stable and can be used as release modifiers in various dosage form [40]. Chirauli nut gum was isolated from the bark of *Buchanania cochinchinesis* (Family: Anacardiaceae) and was used as a release modifier for the preparation of Diclofenac sodium spheroids, using the extrusion spherization technique [41].

**Root:** The rhizome of *Buchanania lanzan* finds an important place in indigenous medicine as an expectorant, diuretic and carminative. It is also found to have anticancer, antihypertensive, larvicidal and antidiabetic activities [42-44]. *In vitro* antioxidant activity of methanolic extract of

root of *Buchanania lanzan* Spreng. exhibited strongest radical scavenging activity (EC<sub>50</sub> 0.24±0.02) by DPPH method in comparison to two other methods and the solvents as well [45]. Patnaik *et al.* [46] in 2011 have reported that the methanolic extract of roots of *Buchanania lanzan* Spreng. could be a potential source of an analgesic and anti-inflammatory agent. For assessing the analgesic activity, two concentrations of the methanolic extract of root (200 and 400 mg/kg orally) were taken for acetic acid-induced writhing model and hot plate reaction time model in Swiss albino mice. Analgesia produced by 400 mg/kg was more significant (p<0.01) when compared with that of 200 mg/kg. Similarly, 400 mg/kg significantly reduced the edema formation of rat paw at 1 and 3 hr. after carrageenin injection (p<0.05 and p<0.01 respectively) when tested for anti-inflammatory activity. The antiulcer activity of ethanolic extract of roots of *Buchanania lanzan* Spreng. was investigated by Kodati *et al.* adopting two experimental models i.e. ethanol induced gastric ulcer and pylorus ligation induced gastric ulcer. Ulcer index parameter was used for the evaluation of antiulcer activity since ulcer formation is directly related to factors such as reduction in gastric volume, decrease in free and total acidity. They found that the oral administration of root's extract at 200-400 mg/kg in pylorus ligation inhibited the total ulcer index by 60.4±2.9 to 69.5±3.7 percent in dose dependent manner as compared with control [47]. Roots of *Buchanania lanzan* Spreng. contain tannins, saponins and flavonoids and the presence of tannins in *Buchanania lanzan* probably contributed to its antidiarrheal activity. Castor oil induced diarrheal test was used to assess the antidiarrheal activity and the gastrointestinal tract transit of charcoal meal test was used to assess the antipropulsive activity of the alcoholic extract of *Buchanania lanzan* Spreng. roots. The study revealed that the extract significantly delayed the onset of diarrhea induced by castor oil and also reduced the number of animals exhibiting diarrhea [48]. Methanolic extract of root of *Buchanania lanzan* Spreng. displayed significant wound healing activity in excision model as also antimicrobial activity against Gram positive (*Staphylococcus aureus* MTCC 96 and *Bacillus subtilis* MTCC 441) and Gram negative (*Escherichia coli* MTCC 2939 and *Pseudomonas aeruginosa* MTCC 2453) bacteria. It was also able to reduce biofilm formation and cause disruption to preformed biofilms in a manner similar to ciprofloxacin [49].

## CONCLUSION

Medicinal plants are integral and indispensable part of the traditional system of medicine practiced

worldwide because of their economical viability, easy accessibility and centuries old experience. As Nature's gift, these are considered to be biocompatible, environment friendly, non-toxic, much cheaper and quite freely available in comparison to synthetic substances. Besides, these are amongst the richest renewable source of biopolymers, having enormous potentials for use and application in numerous fields like foods, cosmetics, pharmaceuticals and host of other industries. With a view to augmenting their sustainable production, conservation, livelihood security and fullest utilization, proper research support is an urgent requirement for addressing the problems related to scientific tapping, harvesting, collection, processing (drying, grading, handling/storage), value addition and upgrading the product quality. This species has a high socio-economic value for providing livelihood to the tribal population of the area besides possessing enormous potentials as commercial horticulture species. Unfortunately due to over-exploitation and indiscriminate harvesting (lopping and cutting), considerable reduction in the population of *Buchanania lanzan* has been recorded in the recent

past, leading to very severe threat to its extinction, which calls for urgent conservation efforts at all levels. The production of this economically important forest tree species is further threatened by insect pests also. In this background, there is compelling need for developing a suitable technology facilitating easy multiplication, regeneration and conservation of the species, simultaneously imparting and disseminating proper knowledge and education to the tribal population for stopping the practice of destructive harvesting and spreading sufficient awareness regarding collection of ripe fruits at appropriate timings.

#### ACKNOWLEDGEMENTS

The authors earnestly express their immense gratitude to Dr. R. Ramani, Director, Indian Institute of Natural Resins & Gums, Namkum, Ranchi (Jharkhand), India for his continuous encouragement and support. They also thankfully acknowledge the financial assistance from Network Project on 'Harvesting, Processing and Value Addition of Natural Resins and Gums', ICAR, New Delhi.

Table 1: Botanical origin and vernacular names of *Buchanania lanzan*

Botanical origin	Vernacular names
Kingdom: Plantae	Bengali: Chironji
Order: Spindales	Hindi: Achar, Char, Baruda, Priyala
Family: Anacardiaceae	Gujarati & Marathi: Charoli, Pyalchar
Sub-family: Anacardioideae	Kannada: Charpoppu
Genus: <i>Buchanania</i>	Malyalam: Mungapper
Species: <i>Buchanania lanzan</i> Spreng.	Oriya: Charu
	Sanskrit: Priyalam, Char, Rajadana, Dhanu.
	Tamil: Morala
	Telugu: Saarachettu, Morichettu.

#### REFERENCES

- Kirtikar KR, Basu BD. Indian Medicinal Plants: Lalit Mohan Basu, 2nd ed., Vol. III, Allahabad, 1935.
- Anon. Wealth of India, A dictionary of Indian raw materials and Indian products, B revised ed., Vol. 2: Publication and Information Directorate CSIR: New Delhi, 1952.
- Pandey GP. Effects of gaseous hydrogen fluoride on the leaves of *Terminalia tomentosa* and *Buchanania lanzan* trees. J Environ Pollution (Series A) 1985; 37:323-334.
- Singh P et al. Preservation of *Buchanania lanzan* Spreng. seeds by *Ocimum canum* Sims essential oil. Annals Plant Protec Sci 2011; 19(2):407-410.
- Sharma VK et al. Phosphate uptake and distribution of the seedling of *Buchanania lanzan*. Tropical Ecol 1970; 11(2):231-232.
- Choubey A et al. Some aspects of germination studies in *Buchanania lanzan* Spreng. seeds. J Tropical Forestry 1997; 13(11):65-73.
- Shukla SK, Solanki KR. Studies on seed germination, plant survival and growth of chironjee (*Buchanania lanzan* Spreng.). J Tropical Forestry 2000;16:44-49.
- Sharma A. Scientific harvesting for quality seed collection of *Buchanania lanzan* Spreng. for its conservation and sustainable management---case study of Chhindwara, Madhya Pradesh, India. Int J Bio-Sci Bio-Tech 2012; 4(1):65-74.
- Kumar J et al. Chironji nut (*Buchanania lanzan*) processing, present practices and scope. Ind J Trad Knowledge 2012; 11(1):202-204.
- Malik SK et al. Genetic resources of chironji (*Buchanania lanzan* Spreng.): a socio-economic important tree species of central Indian population. Genetic Resource Crop Evaluation 2012; 59:615-623.
- Chauhan PS et al. Chironji: a promising tree fruits of dry subtropics. Hort Flora Res Spectrum 2012; 1(3):375-379.
- Bhasin GD, Roonwal ML, Singh B. A list of insect pests of forest plants in India and the adjacent countries. Part 3. Indian Forest Bulletin (NS), Manager of Publication, Govt. of India, New Delhi, 1958; 17:1-11.
- Meshram PB. Tropical forest insects and their control measures, Khanna Bandhu, 7 Tilak Road, Dehradun, 2005; pp. 1-262.

14. Meshram PB, Soni KK. Application of delivery methods for fungal pathogens and insecticides against chironji (*Buchanania lanzan*) stem borer, *Plocaederus obesus* Gahn. Asian J Experiment Biol Sci 2011; 2(1):53-57.
15. Meshram PB, Soni KK. Impact of pest problems and management in *Buchanania lanzan* in Central India. Academic J Entomol 2011; 4(1):30-33.
16. Sengupta A, Roychoudhury SK. Triglyceride composition of *Buchanania lanzan* seed oil. J Sci Food and Agri 1977; 28(5):463-468.
17. Hemavathy J, Prabhakar JV. Lipid composition of chironji (*Buchanania lanzan*) kernel. J Food Composition Anal 1988; 1(4):366-370.
18. Warokar AS et al. Anti-inflammatory and antioxidant activities of methanolic extract of *Buchanania lanzan* Kernel. Ind J Pharmaceu Education Res 2010; 44(4):363-368.
19. Puri A et al. Immunostimulant activity of dry fruits and plant materials used in Indian traditional medical system for mothers after child birth and invalids. J Ethnopharmacol 2000; 71:89-92.
20. Mishra N et al. Study on antioxidant activity of common dry fruits. Food Chem Toxicol 2010; 48:3316-3320.
21. Tyagi N. A novel, potent, bio-film former from the seeds of *Buchanania lanzan* for formulating tobramycin occusers. Int J PharmTech Res 2012; 4(1):422-426.
22. Tyagi N, Madhav NVS. Formulation and evaluation of zidovudine micro emulsion using a novel biopolymer from the seeds of *Buchanania lanzan*. Int J Biopharmaceutics 2012; 3(1):40-43.
23. Rai YC. *Buchanania lanzan* Spreng.----studies on methods of propagation and estimation of fruit yield. Ind Forester 1982; 108(7):501-511.
24. Chitra V et al. Wound healing activity of alcoholic extract of *Buchanania lanzan* in Albino rats. Int J Chem Tech Res 2009; 1(4):1026-1031.
25. Arya R et al. Myricetin 3'-rhamnoside-3-galactoside from *Buchanania lanzan* (Anacardiaceae). Phytochem 1992; 31(7):2569-2570.
26. Mehta SK et al. Preliminary phytochemical investigation on leaves of *Buchanania lanzan* (chironji). Int J Pharmaceu Sci Review Res 2010; 3(2):55-59.
27. Mehta SK et al. Isolation and phytochemical investigation on leaves of *Buchanania lanzan* (chironji). Annals Biol Res 2011; 2(3):469-473.
28. Arya R et al. Flavonoids from the leaves of *Buchanania lanzan* (Anacardiaceae). J Ind Chem Soc 1988; 65:882.
29. Mehrotra S, Mitra R. Pharmacognostical studies on Priyal----*Buchanania lanzan* Spreng. leaf. Bulletin Medico Ethno Bot Res 1981; 1(3):338-352.
30. Mitra R, Mehrotra S. Pharmacognostic studies of 'Priyal'----*Buchanania lanzan* Spreng. stem bark. Herba Hungarica 1981; 20:15-32.
31. Mitra R, Mehrotra S. Pharmacognostic studies on priyal: fruits and seeds of *Buchanania lanzan* Spreng. Bulletin Bot Survey India 1982; 22(1-4):68-76.
32. Verma KS et al. Phytochemical screening and antibacterial activity of *Buchanania lanzan* Spreng. J Phytol Res 2010; 23(2):339-342.
33. Manjunath JR, Mithun NM. Antibacterial and antifungal activity of the *Buchanania lanzan* leaves. Ethnopharmacol, Article ID-"Inventi:ep/403/11" 2011.
34. Mehta SK et al. Adaptogenic activity of methanolic extract of *Buchanania lanzan* leaves an experimental study in rat model. Der Pharmacia Sinica 2011; 2(3):107-112.
35. Sushma N et al. Antidiabetic, antihyperlipidemic and antioxidant activities of *Buchanania lanzan* Spreng. methanol leaf extract in streptozotocin-induced types I and II diabetic rats. Tropical J Pharmaceu Res 2013; 12(2):221-226.
36. Chaudhary US et al. Effect of water extract of the bark of *Buchanania lanzan* Linn. on behaviour and chromatophores of a fresh water fish, Labeorohita. J Environment Biol 2001; 22(3):229-231.
37. Shailasree S et al. Curative properties of *Buchanania lanzan*: As evaluated by its anti-oxidant, anti-inflammatory and DNA protective properties. J Nat Pharmaceuticals 2012; 3(2):71-77.
38. Jain R, Jain SK. Effect of *Buchanania lanzan* Spreng. bark extract on cyclophosphamide induced genotoxicity and oxidative stress in mice. Asian Pacific J Tropical Med 2012; 5(3):187-191.
39. Mehta SK et al. Comparative anti-oxidant activity studies of *Buchanania lanzan* methanolic extract. Biomed Pharmacol J 2013; 2(2):441-444.
40. Bothara SB, Singh S. Thermal studies on natural polysaccharide. Asian Pacific J Tropical Med 2012; S1031-S1035.
41. Gaikwad NB et al. Development of controlled release spheroids using *Buchanania cochinchinesis* gum. J Excipients Food Chemicals 2013; 4(1):4-11.
42. Achuthan CR, Jose P. Hypolipidemic effect of *Alpinia galangal* (Rasna) and *Buchanania lanzan* (Kachoori). Ind J Clinical Biochem 1997; 12(1):55-58.
43. Xue Y, Chen H. Study on the anticarcinogenic effect of three compounds of *Buchanania lanzan* L. Xei Shen Yan Jiu 2002; 31(4):247-248, 251.
44. Othman R et al. Vaso relaxant effect of *Buchanania lanzan* on smooth muscles of rat aorta. Planta Med 2002; 68(7):655-657.
45. Patra KC et al. Anti-oxidant activity of *Buchanania lanzan* Spreng. Family Anacardiaceae. Pharmacologyonline 2011; 1:733-739
46. Patnaik AK et al. Analgesic and anti-inflammatory activities of *Buchanania lanzan* Spreng. roots. Res J Pharmaceu, Biol Chem Sci 2011; 2(1):419-425.
47. Kodati D et al. Antilucer activity of ethanolic extract of *Buchanania lanzan* Spreng. roots. Annals Biol Res 2010; 1(4):234-239.
48. Kodati D et al. Antidiarrhoeal activity of alcoholic extract of *Buchanania lanzan* Spreng. roots. Pharmacologyonline 2010; 3:720-726.
49. Pattnaik A et al. Pharmacological studies on *Buchanania lanzan* Spreng.----A focus on wound healing with particular reference to anti-biofilm properties. Asian Pacific J Tropical Med 2013; 3(12):967-974.